



Rapid Detection of Norwalk-Like Viruses (NLVs)

Satiya Wati
B. App. Sc (Hons)

Infectious Diseases Laboratories
Institute of Medical and Veterinary Science
Adelaide
South Australia

Department of Microbiology and Immunology
University of Adelaide
Adelaide
South Australia

***A thesis submitted to the University of Adelaide in fulfilment of the requirements for
the degree of Master of Science***

July, 1999

Contents

Abstract	vii
Declaration	x
Acknowledgements	xi
Abbreviations	xii
 CHAPTER 1. Introduction	
1.1 Review of gastroenteritis causing agents	1
1.1.1 Bacterial and protozoal gastroenteritis	1
1.1.2 Viral gastroenteritis	2
1.1.2.1 Rotavirus (<i>Reoviridae</i>).....	3
1.1.2.2 Adenovirus (<i>Adenoviridae</i>)	5
1.1.2.3 Astrovirus (<i>Astroviridae</i>)	5
1.1.2.4 Sapporo-like viruses (SLVs) and Norwalk-like viruses (NLVs) (<i>Caliciviridae</i>).....	6
 1.2 History – Discovery of Norwalk-like viruses	 7
 1.3 Significance of NLVs in relation to bacterial and other viral gastroenteritis	 8
 1.4 Classification of NLV.....	 9
1.4.1 Genomic organisation of NLVs	10
1.4.2 Comparison of NLV and SLV genomic organisation	11
1.4.3 Antigenic diversity of NLVs	12
 1.5 Ultra structure of NLV	 13
 1.6 Physicochemical properties of NLVs	 14

1.7	Pathogenesis	15
1.8	Clinical Presentation of NLV infection	16
1.9	Immunity	17
1.10	Epidemiology	18
1.10.1	Transmission	19
1.10.2	Incidence of NLV in Australia	21
1.11	Diagnosis	22
1.11.1	Electron microscopy	22
1.11.2	Immunoassays using human convalescent sera and antigen purified from faeces	23
1.11.3	New approaches to rapid laboratory diagnosis of NLV infection	23
1.11.3.1	Recombinant NLV enzyme immunoassay	24
1.11.3.2	NLV RT-PCR	25
1.12	Project Aims.....	25

CHAPTER 2. General Materials and Methods

2.1	Commonly used reagents	26
2.2	Methods	30
2.2.1	Maintenance of SF21 cells	30
2.2.2	Infection of SF21 cells with recombinant baculovirus	30
2.2.3	Harvest of infected SF21 cells	31
2.2.4	Quantitation of virus particles	31
2.2.5	Electron Microscopy (EM)	32
2.2.6	Density of CAM VLPs	32
2.2.7	Fractionation of antisera with ammonium sulphate	32
2.2.8	Detection of CAM VLP proteins	33

2.2.8.1 Separation of proteins in 10% SDS PAGE gels	33
2.2.8.2 Western blot and detection of CAM VLP proteins	33
2.2.9 Hirt extraction of plasmid DNA from recombinant baculovirus	34
2.2.10 RNA extraction using the Boom method	34
2.2.11 Chomczynski and Sacchi (CS) method	35
2.2.11.1 RNA extraction using the Chomczynski and Sacchi (CS) method	35
2.2.11.2 RNA extraction using the modified Chomczynski and Sacchi (CS) method	36
2.2.12 DNaseI treatment of extracted RNA	36
2.2.13 RNA electrophoresis on Northern gel	37
2.2.14 Poliovirus culture	37
2.2.15 Plaque assay for determining poliovirus plaque forming units (pfu)	38
2.2.16 Negative controls used in EIA and all RT-PCR	38
2.2.17 Processing of faecal specimens to be tested by EIA or RT-PCR	38
2.2.18 Titan™ one tube RT-PCR system	40
2.2.19 RT-PCR of poliovirus using Corbett Thermocycler	40
2.2.20 RT-PCR of rotavirus using Corbett Thermocycler	40
2.2.21 RT-PCR of NLV using Corbett Thermocycler	41
2.2.22 Detection of amplicons	41
2.2.22.1 Electrophoresis of PCR products	41
2.2.22.2 Digoxigenin (DIG) ELISA	41
2.2.23 Sensitivity of RT-PCR amplification	42
2.2.23.1 Competent <i>Escherichia coli</i> (<i>E.coli</i>) strain DH5α	42
2.2.23.2 Plasmid vectors	43
2.2.23.3 Cloning of the 205bp NLV cDNA products into plasmid DNA	43
2.2.23.4 DNA ligation	44
2.2.23.5 Transformation of plasmid into bacteria	44
2.2.23.6 Selection of colonies	45
2.2.23.7 Confirmation of 205bp NLV cDNA in recombinant plasmid DNA	45
2.2.23.8 <i>In vitro</i> transcription	45
2.2.23.9 Quality and quantity of RNA	46
2.2.24 Purification of DNA for sequencing	46
2.2.25 Sequencing analysis	47

2.2.26 Restriction enzyme digestions	47
2.2.27 Phenol extraction and ethanol precipitation of nucleic acids.....	47
2.2.28 List of suppliers	48

CHAPTER 3. Development of a NLV (Camberwell) Enzyme Immunoassay

3.1 Introduction	49
3.1.1 Aims	50
3.1.2 Baculovirus expression system	50
3.1.3 Baculovirus expression of NLV virus-like particles (VLPs)	50
3.2 Materials and Methods	51
3.2.1 Recombinant Baculovirus containing ORF2 of CAM and antisera for detection.....	51
3.2.2 Verification of CAM ORF2 cDNA in clone obtained from Monash University	51
3.2.3 Expression of CAM VLPs	52
3.2.4 Time course experiments to determine optimal day of harvest of CAM VLPs and distribution of these VLPs (cell or supernatant)	53
3.2.5 Production of Recombinant CAM VLPs in Cell Factories	53
3.2.6 Purification of CAM VLPs.....	53
3.2.7 Quantitation of purified CAM VLPs.....	54
3.2.8 Production of anti-CAM serum in rabbits and guinea pig	54
3.2.9 Immunoreactivity of purified CAM VLPs to CAM VLP antisera produced in rabbits and guinea pigs	55
3.2.10 Dilution of purified recombinant Camberwell capsid protein in 10% faecal suspension..	55
3.2.11 EIA to detect CAM antigen	55
3.2.12 Determination of EIA cutoff value.....	56
3.3 Results.....	57
3.3.1 Verification of CAM ORF2 sequence in recombinant Baculovirus.....	57
3.3.2 Expression of recombinant CAM VLPs	57
3.3.3 Purification of CAM VLPs	57

3.3.4	Density of CAM VLPs in Caesium chloride	58
3.3.5	Electron microscopy.....	58
3.3.6	Immunoreactivity of purified CAM VLPs to CAM VLP antisera produced in rabbits and guinea pigs.....	59
3.3.7	Optimization of Cam VLP antisera for CAM EIA	59
3.3.8	Sensitivity of EIA	59
3.3.9	Specificity of CAM EIA	60
3.4	Discussion.....	60

CHAPTER 4. Development of RT-PCR for detection of NLV RNA

4.1	Introduction.....	63
4.1.1	Reverse transcription polymerase chain reaction (RT-PCR)	63
4.1.2	Problems associated with NLV RT-PCR	63
4.1.3	Aims	65
4.1.4	Titan™ One Tube RT-PCR System (Boehringer Mannheim)	66
4.1.5	RNA extraction methods	66
4.2	Materials and Methods	67
4.2.1	Comparison of in-house RNA extraction methods from A549 tissue culture cells and faecal suspensions	67
4.2.2	Development of RT-PCR for poliovirus, rotavirus and NLV	67
4.2.3	Sensitivity of RT-PCR in terms of poliovirus plaque forming unit (pfu)	68
4.2.4	Sensitivity of RT-PCR as judged by end point titration of rotavirus	68
4.2.5	Testing of routine specimens for rotaviral RNA by RT-PCR	69
4.2.6	Sensitivity of amplification of NLV RNA and cDNA	69
4.2.6.1	Determination of level of sensitivity of Titan RT-PCR in terms of copy numbers	69
4.2.6.2	Determination of level of sensitivity of Titan PCR without the RT step	70
4.2.6.3	Determination of level of sensitivity of PCR using Taq polymerase	70

4.3	Results	71
4.3.1	Extraction of total RNA from cells and faecal specimens	71
4.3.2	DNaseI treatment of RNA used for amplification	72
4.3.3	RT-PCR of poliovirus RNA	72
4.3.4	Sensitivity of RT-PCR in terms of poliovirus plaque forming units	73
4.3.5	RT-PCR of rotavirus RNA	73
4.3.6	Routine Specimens tested for rotaviral RNA by RT-PCR	74
4.3.7	Sensitivity of RT-PCR as judged by endpoint titration of rotavirus faecal samples	74
4.3.8	Sensitivity of RT-PCR as judged by endpoint titration of rotavirus in cell culture fluid ..	75
4.3.9	Extraction of NLV RNA and optimization of NLV RT-PCR	75
4.3.10	Determination of the sensitivity of Titan RT-PCR	76
4.3.11	Sensitivity of PCR for DNA alone	76

4.4	Discussion	77
------------	-------------------------	-----------

CHAPTER 5. Testing for NLV faecal samples by CAM EIA and RT-PCR

5.1	Introduction	81
5.1.1	Aims	81
5.2	Materials and Methods	81
5.2.1	Gastroenteritis outbreak specimens in Adelaide (August to December, 1996)	81
5.2.2	Further optimization of NLV RT-PCR to increase sensitivity	82
5.2.3	Comparison of commercial RNA extraction methods to increase sensitivity of RT-PCR	82
5.2.4	Concentration of NLV by Polyethylene glycol MW 8000 precipitation (PEG ppt) and Immuno precipitation (Immuno ppt)	83
5.2.4.1	Concentration of NLV in 41 specimens (Adelaide NLV outbreak) which were negative by RT-PCR	83
5.2.5	Controls for RT-PCR and Nested PCR	84
5.2.6	Nested PCR	84
5.2.6.1	Nested PCR of NLV using Corbett thermocycler	84
5.2.6.2	Sensitivity of nested PCR	84

5.2.6.3 Further amplifications by nested PCR of specimens which were negative for NLV RNA by RT-PCR	85
5.2.7 NLV nucleotide sequences of specimens from the 1996 Adelaide gastroenteritis outbreak	86
5.3 Results	87
5.3.1 Detection of NLV by CAM EIA in stored specimens from 1996 Adelaide gastroenteritis outbreak	87
5.3.2 Detection of NLV RNA by RT-PCR in the outbreak specimens	87
5.3.3 Attempts to increase sensitivity of RT-PCR	88
5.3.3.1 Comparison of two commercial RNA extraction methods to the Boom method to increase sensitivity of RT-PCR	88
5.3.3.2 Comparison of two different concentration methods to increase sensitivity of NLV RT-PCR – precipitation by specific antibodies (Immuno ppt) or with polyethylene glycol (PEG ppt)	88
5.3.3.3 Repeat of RT-PCR on PEG ppt and Qiagen extracted RNA of the 41 first round PCR NLV negative specimens	89
5.3.3.4 Nested PCR	90
5.3.4 Summary of CAM EIA, RT-PCR and nested PCR results on the 65 Adelaide outbreak specimens	91
5.3.5 Inhibition of RT-PCR Enzymes	92
5.3.6 Sequencing	92
5.3.6.1 Sequencing of 205bp RT-PCR and 109bp nested PCR product	92
5.3.6.2 Sequencing of 1443bp nucleotides from the ORF1 region	93
5.3.7 Testing of routine specimens sent to the IMVS for gastroenteritis investigation by CAM EIA and NLV RT-PCR	94
5.4 Discussion	95
CHAPTER 6. General discussion and future work.....	102
References	106

Abstract

NLVs (Norwalk-like viruses) are members of the caliciviridae family and are common causes of epidemic gastroenteritis characterized by short (24-48 hours) incubation and duration. They are currently under diagnosed because of limited availability of convenient diagnostic tests e.g. cell culture systems are not available for this group of viruses. Recent advances including the cloning and sequencing of several strains of this virus has enabled the detection of NLVs in faecal samples. This study describes the development of an RT-PCR and an enzyme immunoassay for the detection of an Australian NLV group 2b virus (Camberwell) and related viruses and their application to clinical samples.

A recombinant baculovirus containing the complete Camberwell capsid protein was provided by Assoc. Prof. Peter Wright (Monash University). Virus-like particles (VLPs) produced by this construct were purified and used to produce hyperimmune rabbit and guinea pig antisera for use in an antigen capture enzyme immunoassay (EIA) with a sensitivity of 0.1ng of purified VLPs. No cross-reaction was observed when a range of gastroenteritis causing bacteria and viruses were used to test the CAM EIA. When applied to a clinical outbreak 31/65 (48%) specimens were found to be EIA positive. In contrast 1000 consecutive routine specimens submitted to our laboratory were EIA negative.

A one tube RT-PCR was also developed for NLV Group 2 (Camberwell-like) viruses. Due to the non-availability of known NLV positive specimens in preliminary experiments, stock cultures of poliovirus and rotavirus positive (by EIA) faecal samples were used for initial optimization of the RT-PCR. The sensitivity of RT-PCR for the three viruses was determined to be 16 pfu for

poliovirus, *ca* 2.5×10^2 viral particles/ml for rotavirus and 2×10^4 copies of *in vitro* transcribed NLV RNA. However, when the outbreak faeces were tested only 24 of the 65 (37%) outbreak specimens were positive on first round NLV RT-PCR.

Further optimization including an improved faecal RNA extraction method (Qiagen RNeasy Plant RNA kit) and concentration of NLVs by Immuno and PEG precipitation methods all improved the sensitivity of the first round RT-PCR but failed to find any additional positive outbreak specimens. However, a nested PCR was found to increase the sensitivity of NLV RT-PCR by 100 fold *in vitro* and found an additional 6 EIA positive specimens to be nested PCR positive. None of the EIA negative specimens became positive. Sequencing of all RT-PCR and nested PCR products showed that they were all identical and showed 96% homology to the Camberwell strain.

This study suggests that NLV antigen capture EIA and RT-PCR assays may be of similar sensitivity and can provide complementary information during the investigation of outbreaks. The development of EIAs for representative strains of other NLV (and Sapporo Like Viruses) genogroups may therefore permit the production of clinically useful diagnostic EIAs. Further development of the RT-PCR assays will require the use of consensus primers for each of the NLV groups 1, 2 and SLV viruses.

Declaration

This thesis contains no material which has been accepted for the award of any other degree or diploma and to the best of my knowledge and belief, the thesis contains no material previously published or written by another person, except where due reference is made in the text. In accordance with the University of Adelaide regulations, I give my consent to this thesis being made available for photocopying and loan if accepted for the award of the degree.

Satiya Wati

Acknowledgments

Many thanks to Dr Tuck Weng Kok and Dr Geoff Higgins for giving me the opportunity to study in the Viral Detection Laboratory and for their supervision.

I would also like to acknowledge the collaboration and helpful discussions with Assoc. Prof. Peter Wright (Monash University, Melbourne) and his staff as well as the provision of the CAM clones, antisera, positive Camberwell specimens and Australian NLV isolate sequences.

Thank you to all the team of the Virus Detection Laboratory for their kindness, help and availability. Special thanks to Adriana Sharley for the graphic work, Sonia Schepetiuk for advice and discussions, Lyn Payne and Sharon Blake for proof-reading and all the past and present staff of VDL for the entertainment during the long working hours.

Special gratitude to fellow peers, John Karlis, Budimen Bela and Adam Davis for the technical advice and continual assistance during the course of my degree. Thanks also to Helen Hocking and Julie Calvert for assistance in obtaining reagents and also encouragement and friendship during my studies.

Thank you to Pete Grant for the photographic work and Rodney Ratcliff for help with alignment of sequences.

Finally I would like to thank my family for all their support and in particular my sister Sanyogita for the generous donation of a computer.

Abbreviations

<i>ca</i>	approximately
e.g.	for example
i.e.	that is
>	greater than
°C	degrees Celsius
bp	base pair
CAM	Camberwell
cDNA	complementary DNA
dATP	2'-deoxyadenosine 5'-triphosphate
DDW	double distilled water
dCTP	2'-deoxycytidine 5'-triphosphate
dGTP	2'-deoxyguanosine 5'-triphosphate
DIG	digoxigenin
DNA	deoxyribonucleic acid
DNaseI	deoxyribonuclease-1
dNTP	2'-deoxynucleoside 5'-triphosphate
DTT	dithiothreitol
dTTP	2'-deoxythymidine 5'-triphosphate
EIA	enzyme immunoassay
EDTA	ethylenediaminetetraacetic acid
EM	electron microscopy
FBS	fetal bovine serum
g (units)	gravity force
IEM	Immune electron microscopy
kb	kilobase pair
kDA	kilodalton
LB	Luria-Bertani medium
µg	microgram
µl	microlitre
µM	micromole per litre (micromolar)
mg	milligram

min(s)	minute(s)
ml	millilitre
mM	millimole per litre (millimolar)
MOPS	3-[N-morpholino]propanesulfonic acid
MW	molecular weight
ng	nanogram
nm	nanometre
NLV	Norwalk-like virus
NV	Norwalk virus
OD	optical density
ORF	open reading frame
PAGE	polyacrylamide gel electrophoresis
pBS	plasmid Bluescript (SK)
PCR	polymerase chain reaction
PEG	polyethylene glycol
p.i.	post infection
ppm	parts per million
ppt	precipitation
RIA	radioimmunoassay
RNA	ribonucleic acid
RNase A	ribonuclease A
rNTP	ribonucleoside 5'-triphosphate
rRNA	ribosomal RNA
rpm	revolutions per minute
RT	reverse transcriptase
SDS	sodium dodecyl sulfate
SLV	Sapporo-like virus
SSC	standard saline citrate
TCA	trichloroacetic acid
UTP	uridine 5'-triphosphate
VLPs	virus-like particles
X-gal	5-bromo-4-chloro-3-indolyl- β -D-galactoside